

Investigation into Rising *Vibrio* Populations in Eastern Coast Oysters

Eric Grigg, Donna L. Rhoads, M.Sc.

Introduction

Vibrio parahaemolyticus is a gram-negative, rod-shaped, and halophilic bacteria that is the leading cause of seafood-borne bacterial gastroenteritis. The gastroenteritis is caused when *Vibrio parahaemolyticus* is consumed with raw oysters and various other forms of seafood. The gastroenteritis is mainly characterized by watery diarrhea along with nausea, vomiting, abdominal cramps, chills, and fever.¹ *Vibrio vulnificus* causes the same gastroenteritis, but can infect the bloodstream of immunocompromised people causing fever, chills, lowered blood pressure, and skin lesions.² Fifty percent of people infected with *Vibrio vulnificus* succumb to the infection.²

Currently, the oysters are harvested and put through a process called depuration to cleanse them of fecal bacteria (coliforms). In this process, oysters or other shellfish are put in a tank with filtered and sterilized seawater under conditions that better facilitate the filter feeding of the oysters. This causes the expulsion of the contents from the oysters. It also enhances the separation of these contaminants from the shellfish and prevents recontamination. The depuration cleans the oysters of faecal bacteria like *Salmonella*, but is less effective at removing other pathogens and viruses in its current design. It is also inconsistent or inefficient at removing biotoxins, and water native bacteria like *Vibrio*.³ A study by the Aquaculture Research Station at Louisiana State University Agricultural Center, while inconclusive, showed evidence that cooling oysters on dry ice has an effect on *Vibrio* populations, however, icing is not a common practice.⁴ In addition, icing the oysters has a high chance of killing them depending their age and size.

Connecticut has seen an increase in *Vibrio* infections over the last two oyster harvesting seasons. As water temperatures increase *Vibrio* bacteria populations expand but the time and rate of expansion is not known. We started the study in late May because gave us the best data. In addition to this, it's also the time when oyster reproduction is at its highest.⁵ This means that there could be a rise in foodborne *Vibrio* infections during the summer (as seen in the past two years). It's estimated that there are about 4500 cases of *Vibrio parahaemolyticus* and many of those not reported.¹ It's also reported that there 95 cases of *Vibrio vulnificus* per year only half of which are culture confirmed.² In addition to this, evidence shows that these numbers are rising.⁶⁻⁷

From 1996 to 2010, *Vibrio* infection reported to COVIS and FoodNet were recorded. According to COVIS, the number of infections annually per 100,000 people increased from 0.09 to 0.28. The number of annual infections per 100,000 people according to FoodNet increased 0.15 to 0.42.⁶

Materials and Methods

Oyster Collection:

16-18 oysters were collected weekly from June 6th to July 31st at oyster beds at the Sound School in New Haven, Connecticut. These oysters were scrubbed with a scrub brush to prevent interior contamination with shell-based organisms. Samples were collected and processed within an hour of collection.

Oyster Processing:

A Waring commercial blender (Model: 51BL32) was placed on a balance and its mass was taken. The oysters were then shucked and their tissue was placed in the blender to create a pooled sample. The blender was massed once more with the tissue inside and the difference between the two measurements was used to determine the mass of the oysters. 100mL of 18ppt artificial seawater was added into the blender and the tissue was blended at full speed for 1 minute. The total volume of the pooled sample was taken by pouring it from the blender into a graduated cylinder.

Vibrio Testing:

50uL aliquots of tissue were taken from the blended sample and placed onto plates containing Chromagar *Vibrio* media. The plates were then placed in a 37°C incubator for 24-48 hours. The plates were then removed from the incubator and the number of colony forming units (CFUs) were counted. This count was used to find the number of CFUs per 50uL. Plates containing more than 300 CFUs were to confluent to count. The presence of *Vibrio* was confirmed using PCR.

DNA Extraction:

Boiling Method – Isolated colonies of *Vibrio parahaemolyticus* and *Vibrio vulnificus* were taken from the plates and placed in separate 1.5mL centrifuge tubes containing 40uL of sterile water. The tubes were then placed in a centrifuge and spun at 13,200 rpm for 5 minutes in order to pellet the bacteria. The supernatant was discarded and 200uL of sterile water was added to each tube. The tubes were then placed in a 100°C water bath for 10 minutes. The tubes were then removed and centrifuged at 13,200 rpm for 2 minutes. The supernatant was used for PCR.
Chelex Method – Isolated colonies of *Vibrio parahaemolyticus* and *Vibrio vulnificus* were taken from the plates and added to separate 1.5mL centrifuge tubes containing 200uL of 10% chelex solution. These tubes were boiled for 10 minutes. The tubes were then centrifuged at 13,200 rpm for 5 minutes. The tubes were removed and the supernatant was transferred to separate 1.5mL centrifuge tubes.

PCR Confirmation:

Tox-130 and Tox-200 primers were used to confirm presence of Vv and VPM-1 and VPM-2 primers were used to confirm presence of Vp. PCR reactions were loaded into Ready-To-Go PCR tubes with 3uL of DNA, 1uL of each primer (forward and reverse), and 21uL of distilled water. The PCR tubes were loaded with each DNA sample being loaded as according the above table. The tubes were then briefly centrifuged. They were then placed in the thermocycler with the following program:

Initial incubation: 94°C for 5 minutes

35 cycles:

94°C for 1 minute

53.4°C for 1 minute

72°C for 1 minute

Final anneal: 72°C for 5 minutes

The PCR products were held at 4°C. 3uL of loading dye was mixed with 7uL of each PCR product. 1.5% agarose gel was prepared and each mixture was loaded into a well. Electrophoresis was run at 100V for 30-45 minutes.



Figure 1. Chromagar *Vibrio* showing presence of *Vibrio parahaemolyticus* (Vp, mauve), *Vibrio vulnificus* (Vv, blue), and *Vibrio algaelyticus* (Va, white).




Figure 4. 1.5% agarose gel showing positive results for Vv and Vp.

Results and Discussion

Oyster Collection:

Our results show an overall increase in *Vibrio* CFUs over the course of the summer. Our first four collections show a constant increase in number of Vv CFUs until our fifth collection. The fifth collection was performed after Hurricane Arthur passed through the Connecticut area around the 4th of July. The hurricane contributed to the drop in Vv numbers by decreasing the salinity of the water and increasing the amount of sediment and runoff from the mainland. A second collection of eight oysters (Collection 5.5) was performed during the same week. The number of CFUs from the second collection dropped slightly from the previous (Collection 5) confirming the large drop in Vv CFUs from week 4 to week 5. The numbers continued to drop for the sixth collection, but they shot up again to a much higher CFU counts for collection 7. The weekend after collection 7, heavy rain came through the area and lowered the *Vibrio* numbers for collection 8.

The trend displayed by the Vp CFUs mimicked the trend set by the Vv, except for the fifth collection which was performed after Hurricane Arthur. The Vp numbers continued to climb while the Vv numbers dropped dramatically. The confirmatory collection showed Vp numbers that were more on par with the trend set by the Vv numbers. The rise in numbers may have arisen from environmental factors such as lowered salinity from the passing storm allowing for the drop in Vv and proliferation of the Vp. The high number could have also arisen from sampling bias.

PCR Confirmation:

The initial Polymerase Chain Reaction confirmed for presence of Vv and Vp. Subsequent runs of the reaction also confirmed presence of *Vibrio*, but were marred by the presence of primer dimers. The primer dimers arise when the primers used stick to each other rather than the template DNA. They stick to each other due to complementary base pairs in each gene sequence. Optimization of the primers is required in order to run an efficient method of detecting the *Vibrio* bacteria.




Figure 2. Number of *Vibrio vulnificus* CFUs per 50uL of tissue.




Figure 3. Number of *Vibrio parahaemolyticus* CFUs per 50uL of tissue

Statistics and Data Analysis

The regression line on the chart of Colony Forming Units (Vv) vs. Weeks has a positive slope indicating an increasing trend. This trend shows that the number of *Vibrio vulnificus* bacteria in Eastern Coast oysters does, in fact, increase over the summer months. This means that the risk of infection with Vv is higher when eating uncooked oysters. The regression line on the chart of Colony Forming Units (Vp) vs Weeks shows the same trend. It displays the same increased risk of infection during the summer months.

The standard deviation of collections from 5 and onward were higher than previous weeks. This is due to the weather conditions that occurred around these times. Week 5's CFU count was lowered due to the passing of Hurricane Arthur. In addition, there was heavy rainfall towards the end of the study.

Future Work

Future studies should focus on *Vibrio* population trends during the Fall, Winter, and Spring months. The populations should fall during the Fall and Winter months due to lowering water temperatures, but they should rise again when Spring begins. The PCR primers need to be optimized or replaced completely and future studies can work to solve the issue with the primer dimers. In addition, a more controlled environment free of weather interference would be beneficial. This would allow us to collect data on *Vibrio* CFUs without the weather disrupting the results.

Conclusion

The study found an increasing trend of both *Vibrio parahaemolyticus* CFUs and *Vibrio vulnificus* CFUs. This shows that the risk of infection with *Vibrio* bacteria via raw oysters rises during the summer months. In addition, heavy rainfall seems to have an effect on *Vibrio* populations. After Hurricane Arthur passed, the CFUs for *Vibrio vulnificus* dropped sharply. This occurred again after heavy rainfall during the eighth week of the study. Given these results, routine surveillance of oyster beds should be instituted. In addition, a more thorough means of cleaning the oysters post-harvest should be applied as the current process, depuration, is inefficient with removing *Vibrio* bacteria.

The PCR confirmed the presence of *Vibrio vulnificus* and *Vibrio parahaemolyticus*, however, some reactions were marred by the presence of primer dimers. These occur when the primers used in the PCR stick to each other rather than to the template DNA. The primers need to be better optimized for efficient use in confirming *Vibrio* species.

Acknowledgements

We would like to thank Rebecca Gratz and the Sound School for the generous use of their space and equipment. We want to thank Gabriel Neil Geist for letting us use his pathology lab and sharing his oyster shucking expertise. We extend a large thanks to the Carrubas for generously financing the SURF Program. We also want to thank Carol Withers and Janice Sanderson for organizing the program and giving me the opportunity to gain the valuable research experience I need for my career. I would also thank fellow SURF Researcher Lawrence Lam and SURF Mentor Donna L. Rhoads for helping me out with the research whenever I needed them to. Lastly, I want to thank the University of New Haven for teaching me and giving me the opportunities and challenges I need to better my academically and personally.

References

- "Vibrio Illness (Vibriosis)." *Centers for Disease Control and Prevention*. Centers for Disease Control and Prevention, 21 Oct. 2013. Web. 05 Mar. 2014. <<http://www.cdc.gov/vibrio/vibriop.html>>.
- "Vibrio Illness (Vibriosis)." *Centers for Disease Control and Prevention*. Centers for Disease Control and Prevention, 21 Oct. 2013. Web. 05 Mar. 2014. <<http://www.cdc.gov/vibrio/vibriov.html>>.
- Lee, Ronald, Alessandro Lovatelli, and Lahsen Ababouch. "Bivalve Depuration: Fundamental and Practical Aspects." *Bivalve Depuration: Fundamental and Practical Aspects*. Food and Agriculture Organization of the United Nations, n.d. Web. 05 Mar. 2014.
- K, Melody, Senevirathne R, James M, Jaykus LA, and Supan J. "Effectiveness of Icing as a Postharvest Treatment for Control of *Vibrio Vulnificus* and *Vibrio Parahaemolyticus* in the Eastern Oyster (*Crassostrea Virginica*)." *Journal of Food Protection* 71.7 (2008): 1475-482. *National Center for Biotechnology Information*. U.S. National Library of Medicine. Web. 05 Mar. 2014.
- "Oysters." *NOAA - Chesapeake Bay Office*. National Oceanic and Atmospheric Administration, n.d. Web. 05 Mar. 2014.
- Newton, Anna, Magdalena Kendall, Duc J. Vugia, Olga L. Henao, and Barbara E. Mahon. "Increasing Rates of Vibriosis in the United States, 1996-2010: Review of Surveillance Data From 2 Systems." *Clinical Infectious Diseases*. Oxford University Press, n.d. Web. 05 Mar. 2014.
- Miller, Ben. "Emerging Pathogens: Vibrio Cases in Oysters Expected to Continue Increasing." *Food Safety News*. N.p., 23 Sept. 2013. Web. 05 Mar. 2014.
- Burgeson, John. "Oysters: A Complicated Economy and Ecology." *Connecticut Post*. Hearst Media Services Connecticut, LLC, 25 May 2013. Web. 10 Mar. 2014.
- "VIBRIO VULNIFICUS." *OLR Research Report*. Connecticut General Assembly, 22 June 2012. Web. 12 Mar. 2014.
- Kim, H. S., D.-M. Kim, G. P. Neupane, Y.-M. Lee, N.-W. Yang, S. J. Jang, S.-I. Jung, K.-H. Park, H.-R. Park, C. S. Lee, and S. H. Lee. "Comparison of Conventional, Nested, and Real-Time PCR Assays for Rapid and Accurate Detection of *Vibrio Vulnificus*." *Journal of Clinical Microbiology* 46.9 (2008): 2992-998. Web.
- Luan, X.-Y., J.-X. Chen, X.-H. Zhang, J.-T. Jia, F.-R. Sun, and Y. Li. "Comparison of Different Primers for Rapid Detection of *Vibrio Parahaemolyticus* Using the Polymerase Chain Reaction." *Letters in Applied Microbiology* 44.3 (2007): 242-47. Web.